

Self-Induced Compactness in Banach Spaces

P.G.CASAZZA AND H.JARCHOW

Introduction

The question which led to the title of this note is the following:

If X is a Banach space and K is a compact subset of X , is it possible to find a compact, or even approximable, operator $v : X \rightarrow X$ such that $K \subset \overline{v(B_X)}$?

This question was first posed by P.G.Dixon [6] in connection with investigating the problem of the existence of approximate identities in certain operator algebras. We shall provide a couple of observations related to the above question and give in particular a negative answer in case of approximable operators.

We shall also provide the first examples of Banach spaces having the approximation property but failing the bounded compact approximation property though all of their duals do even have the metric compact approximation property.

Approximate Identities

A *left approximate identity* (LAI) in a Banach algebra A is a net $(e_i)_{i \in I}$ in A such that $\lim_{i \in I} \|e_i x - x\| = 0$ for each $x \in A$. If there is a $\lambda > 0$ such that in addition $\|e_i\| \leq \lambda$, then $(e_i)_{i \in I}$ is called a λ -*bounded LAI* (λ -BLAI); note that necessarily $\lambda \geq 1$. We say that $(e_i)_{i \in I}$ is a BLAI if it is a λ -BLAI for some λ .

Right approximate identities (RAI's) and (λ) -*bounded RAI's* ((λ) -BRAI's) are defined analogously.

A result which, when seen in the context of operator algebras, is of particular interest for Banach space theory, is P.J.Cohen's [4] factorization theorem:

If the Banach algebra A has a BLAI, then every $z \in A$ can be written as a product $z = xy$ with $x, y \in A$; in addition, y exists in the closed left ideal generated by z and can be chosen so that $\|z - y\| \leq \delta$, where δ was given previously.

See the book [2] of F.F.Bonsall and J.Duncan for details.

An interesting case occurs when A is of the form $\mathcal{A}(X)$ where $\mathcal{A}$ is any Banach operator ideal in the sense of A.Pietsch [13] and X is a Banach space. However, in such generality

approximate identities have hardly been discussed; attention has been focused on the cases where $\mathcal{A}$ is either

$$\mathcal{K},$$

the ideal of all compact operators, or

$$\overline{\mathcal{F}},$$

the ideal of all *approximable operators*, i.e., uniform limits of operators of finite rank operators. Of course,

$$\mathcal{F}$$

will be used to denote the ideal of all finite rank operators between Banach spaces. Dixon's paper [6] is dealing with precisely these ideals, and one of his main results is the following:

Theorem A. *A Banach space X has the λ -BAP [resp. the λ -BCAP] if and only if $\overline{\mathcal{F}}(X)$ [resp. $\mathcal{K}(X)$] has a λ -BLAI.*

Recall that a Banach space X has *AP* (“*approximation property*”) [resp. *CAP* (“*compact approximation property*”)] if, for any compact subset K of X and any $\varepsilon > 0$, there is an operator $u \in \mathcal{F}(X)$ [resp. $u \in \mathcal{K}(X)$] such that $\|ux - x\| < \varepsilon$ for all $x \in K$. If there is a $\lambda (\geq 1)$ such that we always can arrange for $\|u\| \leq \lambda$, then we say that X has λ -BAP [resp. λ -BCAP], with “B” being shorthand for “bounded”, of course. Usually, *BAP* [resp. *BCAP*] is used when we only know that we have λ -BAP [resp. λ -BCAP] for some λ . The case $\lambda = 1$ corresponds to what is usually called *MAP* (“*metric approximation property*”) and *MCAP* (“*metric compact approximation property*”), respectively.

Only recently C.Samuel [16] and N.Grønbæk and G.A.Willis [7] addressed the corresponding problem of existence of RAI's. One of the main results in [7] is the following companion result of the preceding theorem:

Theorem B. *Let X be a Banach space. Then X^* has the λ -BAP if and only if $\overline{\mathcal{F}}(X)$ has a λ -BRAI.*

The situation is less pleasant when $\overline{\mathcal{F}}$ is replaced by $\mathcal{K}$. It is still true that if $\mathcal{K}(X)$ has a λ -BRAI, then X has λ -BCAP. However, the converse fails. As we shall see, there is a Banach space X with AP, but which fails BCAP, while $X^*, X^{**}, \dots$ are all separable and have MCAP. If $\overline{\mathcal{F}}(X) = \mathcal{K}(X)$ has a BRAI, then X^* would have BAP, and so X should have BAP as well – but it doesn't.

To construct our example we need some results from the literature. Recall that a Banach space X has *shrinking* λ -CAP if there is a net $u_\alpha \in \mathcal{K}(X)$, $\|u_\alpha\| \leq \lambda$, (u_α) converges strongly to the identity on X and (u_α^*) converges strongly to the identity on X^* . In his memoir [8], A. Grothendieck showed that a separable dual space with AP has MAP. The corresponding question for CAP is still open. However, an alternative proof of Grothendieck's result due to Lindenstrauss and Tzafriri [11] will work for shrinking CAP (see Cho and Johnson [3]).

Theorem C. *If X^* is separable and has CAP given by w^* -continuous operators, then X and X^* have MCAP.*

It follows that,

Corollary D. *If X^* is separable and X has shrinking CAP, then for every equivalent norm $|\cdot|$ on X , both X and $(X, |\cdot|)^*$ have MCAP.*

We are now ready for:

Theorem 1. *There is a Banach space X which has AP, but fails BCAP, while $X^*, X^{**}, \dots$ are all separable and have MCAP.*

Proof: Let Y be a separable reflexive Banach space with CAP which fails to have AP, see G.Willis [19]. Choose a Banach space Z so that Z^{**} has a basis and $Z^{**}/Z \approx Y$. (This result and the others used here can be found in [11]). Now, $Z^{***} \approx Z^* \oplus Y^*$ fails AP but has shrinking CAP since (1) Z^* has a shrinking basis and (2) Y is reflexive and has CAP. It follows that Y has shrinking CAP. By a construction of Figiel and Johnson ([11], p.42), for each n , there is an equivalent norm $|\cdot|_n$ on Z^{**} so that $(Z^{**}, |\cdot|_n)$ fails n -BAP. But $(Z^{**}, |\cdot|_n)^*$ still has shrinking CAP (being isomorphic to Z^{***}). So by Corollary D, this space and its dual have MCAP. (A similar argument shows, in fact, that all of its duals have MCAP.)

Let

$$X = \left(\sum_{n=1}^{\infty} \oplus (Z^{**}, |\cdot|_n) \right)_{\ell_2}.$$

Then X has AP and fails BAP. So X fails BCAP. (It is easily seen that X has λ -BAP if and only if X has AP and λ -BCAP). The spaces $X^*, X^{**}, \dots$ are all ℓ_2 -sums of Banach spaces having MCAP, and hence have MCAP. This completes the construction. QED

The Properties $(\overline{\mathcal{F}})$ and $(\mathcal{K})$

We shall now concentrate on left approximate units. There is a natural question related to Theorem A: if the same λ works for the BLAI and for the B(C)AP, couldn't it be simultaneously 'eliminated' on both sides? More precisely, is it true that $\overline{\mathcal{F}}(X)$ resp. $\mathcal{K}(X)$ has a LAI if and only if X has AP resp. CAP?

Let $\mathcal{A}$ be $\overline{\mathcal{F}}$ or $\mathcal{K}$. We say that a Banach space X has the *property* $(\mathcal{A})$ if for each compact subset K of X there is an operator $u \in \mathcal{A}(X)$ such that $K \subset u(B_X)$. This is what "self-induced compactness" in the title is referring to. Of course, the concept can be generalized in many directions, but we prefer to stay with the present setup.

The following results are again due to Dixon [6]:

Theorem E. *Let X be a Banach space.*

(a) *If X has AP [CAP], then $\overline{\mathcal{F}}(X)$ [$\mathcal{K}(X)$] has a LAI.*

- (b) If X has property $(\overline{\mathcal{F}})$ $[(\mathcal{K})]$ and if $\overline{\mathcal{F}}(X)$ $[\mathcal{K}(X)]$ has a LAI, then X has AP [CAP].
- (c) If X has BAP [BCAP], then it has property $(\overline{\mathcal{F}})$ $[(\mathcal{K})]$.

All we need to get started is a workable condition which is equivalent to property $(\mathcal{A})$. This is elementary:

Proposition 1. *The following statements about the Banach space X are equivalent:*

- (i) X has property $(\mathcal{A})$.
- (ii) Given $u \in \mathcal{K}(\ell_1, X)$, there are operators $v \in \mathcal{A}(X)$ and $w_n \in \mathcal{L}(\ell_1, X)$ ($n \in \mathbf{N}$) such that $\lim_{n \rightarrow \infty} \|u - vw_n\| = 0$.
- (iii) same as (ii), but $w_n \in \mathcal{K}(\ell_1, X)$.
- (iv) For each $u \in \mathcal{K}(\ell_1, X)$, there are operators $v \in \mathcal{A}(X)$ and $w \in \mathcal{L}(\ell_1, X^{**})$ such that $u = v^{**}w$.
- (v) same as (iv), but $w \in \mathcal{K}(\ell_1, X^{**})$.

Since we are working with (weakly) compact operators $v : X \rightarrow X$, we may and shall consider v^{**} as an operator $X^{**} \rightarrow X$; accordingly we have

$$\overline{v(B_X)} = v^{**}(B_{X^{**}}).$$

Proof. (i) $\Rightarrow$ (ii): Given $u \in \mathcal{K}(\ell_1, X)$, we can find for each $n \in \mathbf{N}$ vectors $x_k \in B_X$ such that $\|vx_k - ue_k\| \leq n^{-1}$ for all $k \in \mathbf{N}$, and then define $w_n : \ell_1 \rightarrow X$ via $w_n e_k = x_k$ for each k .

(ii) $\Rightarrow$ (iii): Any $u \in \mathcal{K}(\ell_1, X)$ can be written as $u = u_1 u_2$ where $u_1 : \ell_1 \rightarrow X$ and $u_2 : \ell_1 \rightarrow \ell_1$ are compact operators.

(iii) $\Rightarrow$ (iv): We may assume that $\|w_n\| \leq 1$ for each n . Let $\mathcal{U}$ be a free ultrafilter on $\mathbf{N}$ and define $w : \ell_1 \rightarrow X^{**}$ by $w(\xi) = \lim_{\mathcal{U}} w_n \xi$, the limit being taken in the weak* topology of X^{**} . This is a bounded linear operator, and $u = v^{**}w$.

(iv) $\Rightarrow$ (v) is obtained as was (ii) $\Rightarrow$ (iii), so we are left with (v) $\Rightarrow$ (i): Let $K \subset X$ be compact. Then $K \subset \overline{\text{conv}} \{x_n : n \in \mathbf{N}\}$ for some null sequence (x_n) in K . The operator $u : \ell_1 \rightarrow X$ defined by $ue_n := x_n$ for each n is compact, so there are $v \in \mathcal{A}(X)$ and $w \in \mathcal{K}(\ell_1, X^{**})$ such that $u = v^{**}w$. We may assume that $\|w\| \leq 1$ so that $K \subset \overline{u(B_{\ell_1})} \subset v^{**}(B_{X^{**}})$. QED

Since ℓ_1 enjoys the lifting property, a Banach space X has property $(\mathcal{A})$ whenever the following applies: no matter how we choose the Banach space Y and the operator $u \in \mathcal{A}(Y, X)$, we can find a quotient Q of X along with operators $v \in \mathcal{A}(Q, X)$ and $w \in \mathcal{L}(Y, Q^{**})$ such that $u = v^{**}w$. It is interesting to note that in case $\mathcal{A} = \mathcal{K}$, the preceding proposition allows to extend this almost to a characterization. We have the following weak version of the Cohen factorization theorem.

Proposition 2. *Let X be a Banach space.*

- (a) If X has property $(\mathcal{K})$ then, given any Banach space Y , every operator $u \in \mathcal{K}(Y, X)$ admits a compact factorization through some quotient Q of X^{**} : there are operators $v \in \mathcal{K}(Q, X)$ and $w \in \mathcal{K}(Y, Q)$ such that $u = vw$.
- (b) Suppose there is, for every Banach space Y and every $u \in \mathcal{K}(Y, X)$, a quotient space Q of X together with operators $v \in \mathcal{K}(Q, X)$ and $w \in \mathcal{K}(Y, Q^{**})$ such that $u = v^{**}w$. Then X has property $(\mathcal{K})$.

Proof. (a) Suppose that X has $(\mathcal{K})$ and let Y and $u \in \mathcal{K}(Y, X)$ be given. Of course we may assume that Y is separable; so we can work with a quotient map $q : \ell_1 \rightarrow Y$. Thanks to Proposition 1 there are operators $v_0 \in \mathcal{K}(X)$ and $w_0 \in \mathcal{K}(\ell_1, X^{**})$ such that $uq = v_0^{**}w_0$. Set $Q = X^{**}/\ker(v_0^{**})$, let $p : X^{**} \rightarrow Q$ be the quotient map, and let $v \in \mathcal{K}(Q, X)$ be such that $vp = v_0^{**}$. As v is injective, there is a $w \in \mathcal{K}(Y, Q)$ such that $wq = pw_0$. Note that $u = vw$.

- (b) Apply the hypothesis to any $\mathcal{K}(\ell_1, X)$ and use the lifting property of ℓ_1 . QED

We do not have a corresponding result for the property $(\overline{\mathcal{F}})$.

We continue by giving a number of immediate consequences of Proposition 1.

If the Banach space X admits a quotient which is isomorphic to ℓ_1 then, by the lifting property of ℓ_1 , this quotient is isomorphic to a complemented subspace of X . So we may state:

Corollary 1. *Any Banach space which admits a quotient isomorphic to ℓ_1 has the property $(\overline{\mathcal{F}})$.*

If X fails CAP then $X \oplus \ell_1$ fails CAP and has ℓ_1 as a quotient space. It follows from Theorem E that for such a space X neither $\mathcal{K}(X)$ nor $\overline{\mathcal{F}}(X)$ can have a LAI. We can easily extend the list of such examples.

Lemma. *Let X, Y and Z be Banach spaces. Suppose that X is a quotient of Y , that Z is a quotient of X , and that every $u \in \mathcal{K}(\ell_1, Y)$ can be written $u = vw$ where $w \in \mathcal{K}(\ell_1, Z)$ and $v \in \mathcal{A}(Z, X)$. Then X has the property $(\mathcal{A})$.*

This is an immediate consequence of the fact that ℓ_1 has the compact lifting property.

Corollary 2. *Let X be a subspace of c_0 or ℓ_p ($1 \leq p < \infty$). Then X^* has property $(\overline{\mathcal{F}})$.*

But X^* may well fail to have CAP in which case again neither $\overline{\mathcal{F}}(X)$ nor $\mathcal{K}(X)$ can have a LAI.

The proof is immediate from the lemma and the fact that every infinite dimensional subspace of c_0 or ℓ_p contains a subspace which is isomorphic to c_0 resp. ℓ_p and complemented in the whole space.

We do not know if every subspace of c_0 or ℓ_p ($1 \leq p < \infty$) has $(\mathcal{K})$ or $(\overline{\mathcal{F}})$. Also, we do not know how $(\mathcal{A})$ behaves with respect to duality.

W.B.Johnson has proved in [10] that there is a separable Banach space, C_1 , such that if Z is any separable Banach space, then Z^* is isometrically isomorphic to a norm-one

complemented subspace of C_1^* . It follows that C_1^* fails CAP but it has the property $(\overline{\mathcal{F}})$ since ℓ_1 is complemented in C_1^* . Consequently, neither $\mathcal{K}(C_1^*)$ nor $\overline{\mathcal{F}}(C_1^*)$ can have a LAI.

On the basis of this it is tempting to conjecture that the property $(\mathcal{A})$ is preserved under the formation of complemented subspaces. The above observation about duals of subspaces of c_0 and ℓ_p would then appear as a consequence of C_1^* 's property of having $(\overline{\mathcal{F}})$; in fact the dual of any separable Banach space would have $(\overline{\mathcal{F}})$. We shall now see that this is not the case.

Let Γ_2 denote the ideal of all Banach space operators which factor through some Hilbert space.

Proposition 3. *Suppose that the Banach space X has the property $(\mathcal{A})$. If $\mathcal{A}(X) \subset \Gamma_2(X)$, then X is isomorphic to a Hilbert space.*

Proof. In fact, Grothendieck's Inequality informs us that $\mathcal{K}(\ell_1, X)$ consists of absolutely summing operators only. By trace duality, however, this can only happen when X is isomorphic to a Hilbert space. QED

It was shown by G. Pisier [14] that $\overline{\mathcal{F}}(X)$ is contained in $\Gamma_2(X)$ whenever X and X^* both have cotype 2, and a few years later, he proved [15] that every cotype 2 space Z embeds into a non-hilbertian space P_Z which, together with its dual, has cotype 2. Such a space necessarily fails AP. Thus:

Corollary 3. *The Pisier spaces P_Z fail to have property $(\overline{\mathcal{F}})$.*

The same is true for all the duals of the spaces P_Z .

But thanks to Corollary 1, the spaces $P_Z \oplus \ell_1$ enjoy property $(\overline{\mathcal{F}})$. Conclusion:

Corollary 4. *The property $(\overline{\mathcal{F}})$ is not preserved when passing to complemented subspaces.*

Another way to obtain this is by using Johnson's universal space C_1^* : if Z is separable, then Pisier's construction leads to a separable space P_Z , so that P_Z^* is complemented in C_1^* .

Actually, the spaces P_Z enjoy even more exotic properties. For example, $\overline{\mathcal{F}}(P_Z)$ coincides with $\mathcal{N}(P_Z)$, the algebra of all *nuclear* operators $u : P_Z \rightarrow P_Z$. We do not know if $\overline{\mathcal{F}}(P_Z) = \mathcal{N}(P_Z)$ can have a LAI; actually, the question of what the meaning of the existence of LAI's in $\mathcal{N}(X)$ is in terms of X doesn't seem to have been investigated. However, it was recently shown by Selivanov [17] that $\mathcal{N}(X)$ has a BLAI if and only if X is finite dimensional; see also Dales and Jarchow in [5].

The arguments employed before also show that $\Gamma_2(P_Z)$ doesn't have a LAI.

Odds and Ends

We start by listing a few open problems:

- (a) Are there Banach spaces failing $(\mathcal{K})$?
- (b) Are there Banach spaces failing $(\mathcal{K})$ such that $\mathcal{K}(X)$ has (doesn't have) a LAI?
- (c) Can any of the algebras $\overline{\mathcal{F}}(P_Z)$ have a LAI?
- (d) When do the algebras $\mathcal{N}(X)$, $\Gamma_2(X)$,..... have a LAI?
- (e) How do the properties $(\mathcal{K})$ and $(\overline{\mathcal{F}})$ behave with respect to duality?
- (f) Regarding $(\mathcal{K})$ and $(\overline{\mathcal{F}})$, what can be said when the underlying Banach space X is a Banach lattice, H^∞ , a C^* -algebra, ?

Here are some further ideas which lead to many more problems.

The property $(\mathcal{A})$ can be generalized as follows. Given a Banach space X , let

$$\mathcal{A}_X$$

be the collection of all Banach spaces Z such that for each compact subset K of X there is an operator $u \in \mathcal{A}(Z, X)$ such that $K \subset \overline{u(B_Z)}$; again $\mathcal{A}$ is $\mathcal{K}$ or $\overline{\mathcal{F}}$.

It seems plausible that investigation of such a concept could be helpful in understanding compactness in general Banach spaces through known characterizations of compactness in e.g. classical spaces. Not much, however, is known, and what is known indicates that the picture will by no means be easy to understand.

There are Banach spaces Z which belong to $\bigcap_X \overline{\mathcal{F}}_X$: ℓ_1 and the duals of Johnson's spaces C_p provide examples.

If X has BAP [resp. BCAP], then X belongs to $\overline{\mathcal{F}}_X$ [resp. $\mathcal{K}_X$], whereas the Pisier spaces P_Z satisfy $P_Z \notin \overline{\mathcal{F}}_{P_Z}$.

These spaces also satisfy $P_Z \notin \mathcal{K}_{P_Z^*}$. In fact, K.John [9] has shown that every compact operator $P_Z \rightarrow P_Z^*$ is nuclear, so that $P_Z \in \mathcal{K}_{P_Z^*}$ would entail $\mathcal{K}(\ell_1, P_Z^*) = \mathcal{N}(\ell_1, P_Z^*)$ which cannot be reconciled with P_Z being infinite-dimensional.

On the other hand, there are Banach spaces X such that $X \in \overline{\mathcal{F}}_{X^*}$: think of $X = \ell_1$, $X = C_p$, $X = \ell_2$,.... Can one characterize such spaces?

By the same type of argument we get that if X is a non-hilbertian cotype 2 space, then $\overline{\mathcal{F}}_X$ cannot contain any Z such that Z^* has cotype 2. In fact, otherwise we would get $\mathcal{K}(\ell_1, X) = \Gamma_2(\ell_1, X)$ (cf. [14]), and this is only possible if X is isomorphic to a Hilbert space. Similar, if X has cotype 2 and if $\mathcal{K}_X$ contains a Banach space of type 2, then X must be hilbertian.

All these example revolve around Hilbert space and amount to ending up with the conclusion that $\mathcal{K}_X$ contains a Hilbert space iff X is isomorphic to Hilbert space.

We claim that, if X is hilbertian, then $\mathcal{K}_X$ contains even all dual Banach spaces. (All spaces are supposed to be infinite dimensional). This can be seen as follows:

Let $u : \ell_1 \rightarrow X$ be a compact operator. Since X is hilbertian, $u : \ell_1 \xrightarrow{u_2} \ell_2 \xrightarrow{u_1} X$, where u_1 and u_2 are compact operators. Let now Z be any Banach space. By a result of S.Bellenot [1] (J.S.Morell and J.R.Retherford [12]) there exists a quotient space Q of Z^*

and compact operators $v_1 : Q \rightarrow X$ and $v_2 : \ell_2 \rightarrow Q$ such that $u_1 = v_1 v_2$. Invoke the compact lifting property of ℓ_1 to finish the proof.

The situation resembles the one encountered in Proposition 2. Question: Can one get a factorization through Z rather than through Z^* ?

REFERENCES

- [1] S.Bellenot: The Schwartz-Hilbert variety.
Mich.Math.J. **22** (1975) 373 -377.
- [2] F.F.Bonsall and J.Duncan: Complete Normed Algebras.
Springer-Verlag 1973.
- [3] C.Cho and W.B.Johnson: A characterization of subspaces X of ℓ^p for which $K(X)$ is an M-ideal in $L(X)$. Proc.AMS **93** (1985) 466-470.
- [4] P.J.Cohen: Factorization in group algebras.
Duke Math.J. **26** (1959) 199-205.
- [5] H.G.Dales and H.Jarchow: Continuity of homomorphisms and derivations from algebras of approximable and nuclear operators.
Math.Proc.Cambr.Phil.Soc., to appear.
- [6] P.G.Dixon: Left approximate identities in algebras of compact operators on Banach spaces.
Proc.Roy.Soc.Edinburgh **104A** (1989)169 -175.
- [7] N.Grønbæk and G.A.Willis: Approximate identities in the Banach algebras of compact operators. Canad.Math.Bull.
- [8] A. Grothendieck: Produits tensoriels topologiques at espaces nucléaires.
Mem.Amer.Math.Soc. **16**, 1955.
- [9] K.John: On the compact non-nuclear problem.
Math.Ann. **287** (1990) 509-514.
- [10] W.B.Johnson: A complementary universal conjugate Banach space and its relation to the approximation problem.
Israel J.Math. **13** (1972) 301-310.
- [11] J.Lindenstrauss and L.Tzafriri; Classical Banach Spaces I, Sequence Spaces.
Springer-Verlag, Erg.Math.Grenzg. **92**, 1977.
- [12] J.S.Morell and J.R.Retherford: p -trivial Banach spaces.
Studia Math. **43** (1972) 1-25.
- [13] A.Pietsch: Operator Ideals.
VEB Deutscher Verlag der Wissenschaften 1978 / North-Holland 1980.
- [14] G.Pisier: Un théorème sur les opérateurs entre espaces de Banach qui se factorisent par un espace de Hilbert.
Ann. École Norm.Sup. **13** (1980) 23-43.
- [15] G.Pisier: Counterexample to a conjecture of Grothendieck.

- Acta Math. **151** (1983) 180-208.
- [16] C.Samuel: Bounded approximate identities in the algebra of compact operators on a Banach space.
Proc.AMS **117** (1993) 1093-1096.
- [17] Y.V.Selivanov: Homological characterizations of the approximation property for Banach spaces.
Glasgow Math.J. **34** (1992) 229-239.
- [18] A.Szankowski: Subspaces without the approximation property.
Israel J.Math. **30** (1978) 123-129.
- [19] G.A.Willis: The compact approximation property does not imply the approximation property.
Studia Math. **103** (1992) 99-108.

P.G.Casazza
Department of Mathematics
University of Missouri-Columbia
Columbia, Mo 65211
USA
e-mail:
Pete@casazza.cs.missouri.edu

H.Jarchow
Mathematisches Institut
Universität Zürich
CH 8057 Zürich
Switzerland
e-mail:
Jarchow@math.unizh.ch